



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

The particles are sliding down inclined planes, inclined at an angle $\beta = \frac{1}{4}\pi$ to the horizon, starting from rest. Let a = side of square, x = distance each particle has moved after a time t . Then the velocity of each is

$$v = \sqrt{(2g \sin \frac{1}{4}\pi \cdot x)} = \sqrt{(gx \cdot 2)} = \sqrt{(ga \cdot 2)} \text{ at } D.$$

Each particle is moving toward the diagonal at a velocity $= v \sin \frac{1}{4}\pi = \frac{1}{2}v \cdot \sqrt{2} = \frac{1}{2}\sqrt{[2\sqrt{(2)gx}]} = \frac{1}{2}\sqrt{[2\sqrt{(2)ga}]}$ at D . Hence, the particles at D approach each other with a velocity $= 2 \times \frac{1}{2}\sqrt{[2\sqrt{(2)ga}]} = \sqrt{[2\sqrt{(2)ga}]}$.

Let e = the coefficient of restitution. Then, since the particles impinge at right angles, they will separate with a velocity $v_1 = e\sqrt{[2\sqrt{(2)ga}]}$, and will ascend DB , DA , respectively, with a velocity $v_2 = e\sqrt{(ga \cdot 2)}$. Each will ascend a distance $= v_2^2 / 2g \sin \frac{1}{4}\pi = v_2^2 / g \cdot \sqrt{2} = e^2 a$.

Therefore, they separate, after impact, a distance $= 2e^2 a \sin \frac{1}{4}\pi = e^2 a \sqrt{2}$. If $e=1$, they return to their starting points.

AVERAGE AND PROBABILITY.

190. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

A line is drawn at random across a regular $2n$ -gon; what is the chance that it crosses parallel sides?

Solution by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue, Philadelphia, Pa.

Let AB , CD be two parallel sides. Through the center of the polygon O draw PQ to represent the direction of the random line.

Let $AO=r$, $\angle AOB = \pi/n = \beta$, $\angle AOP = \theta$.

For all possible positions PQ may be regarded as lying in one quadrant. The actual line parallel to PQ may hold any position within breadth of plane HF (H being the most remote vertex from QP , and F foot of perpendicular from H on QP) for all positions, and AE (A being the nearest vertex to QP and E foot of perpendicular from A to QP) for all favorable positions.

\therefore The chance is $p = AE/HF$; $AE = r \sin \theta$, $HF = r \cos \theta$ for n even; $HF = r \cos(\frac{1}{2}\beta - \theta)$ for n odd. For n even,

$$p = \frac{2n}{\pi} \int_0^{\frac{1}{2}\beta} \tan \theta d\theta = \frac{2n}{\pi} \log(\sec \frac{1}{2}\beta). \quad \therefore p = \frac{2n}{\pi} \log \left(\sec \frac{\pi}{2n} \right).$$

$$\text{For } n \text{ odd, } p = \frac{2n}{\pi} \int_0^{\frac{1}{2}\pi} \frac{\sin \theta}{\cos(\frac{1}{2}\beta - \theta)} d\theta = \frac{n}{\pi} (\beta \sin \frac{1}{2}\beta + 2 \cos \frac{1}{2}\beta \log \cos \frac{1}{2}\beta).$$

$$\therefore p = \frac{n}{\pi} \left[\frac{\pi}{n} \sin \frac{\pi}{2n} + 2 \cos \frac{\pi}{2n} \log \left(\cos \frac{\pi}{2n} \right) \right].$$